

The NIMH global mental health research community and COVID-19



The world faced substantial challenges in meeting the demands for mental health care, even before the emergence of coronavirus disease (COVID-19). With the havoc caused by the pandemic and the impending impact on economies, social structures, and health systems, a global mental health crisis is arising. The Director-General of the UN forecast this situation in a policy brief on May 13, 2020, stating, "The mental health and wellbeing of whole societies have been severely impacted by this crisis and are a priority to be addressed urgently."¹ Three critical actions were recommended: apply a whole-of-society approach to promote, protect, and care for mental health; ensure widespread availability of emergency mental health and psychosocial support; and support recovery from COVID-19 by developing mental health services for the future.

In a field that is chronically underfunded and ignored by most policy makers, especially in low-income and middle-income countries (LMICs), taking action is not easy. Upwards of 90% of people with mental health conditions receive no treatment in LMICs. As the COVID-19 pandemic unfolds, and given the likelihood of even fewer resources dedicated to mental health services, questions arise about whether it will be feasible to implement the UN's recommendations in the coming years.

Can the research community offer any rapid solutions? The National Institute of Mental Health (NIMH) supports a network of global research Hubs designed to address questions that arise as LMICs widely implement sustainable, evidence-based mental health services.² These so-called "NIMH Scale-Up Hubs" are interdisciplinary, with interest and expertise of the people involved in increasing the reach, accessibility, adoption, quality, costs, and effectiveness of mental health services, enhancing collaborative learning and development, building local capacity for implementation research, and establishing relationships with governmental, non-governmental, and community-based stakeholders. We present examples of replicable programmes that can support the UN's recommendations.

Hubs deploy creative strategies to engage entire communities in delivering evidence-based interventions through various platforms across the life course. In

Pakistan, teachers are trained to recognise and manage emotional and behavioural problems in children attending primary and secondary schools through a specially developed online training programme integrated into the teachers' ongoing education platforms.³ Teachers use an interactive chatbot to work with children in classroom settings, and become better equipped to prevent any emotional or behavioural problems from disrupting the children's education and wellbeing. In Uganda, parents are trained to work alongside community health workers to deliver a family group intervention to children with disruptive behaviours.⁴ The intervention builds support for families by providing opportunities for caregivers and children to communicate in safe settings with other families who have shared experiences. In Sierra Leone, evidence-based group interventions for mental health are integrated into youth entrepreneurship programmes that offer war-affected young people opportunities for a livelihood and encourage development of local capacity.⁵ The interventions combine elements from cognitive behavioural and interpersonal therapies and are delivered by lay workers. In India, community-based interventions for suicide prevention are implemented among students and farmers, with community health workers who are trained to identify and refer vulnerable groups to local mental health services.⁶ In Thailand, people involved in community-based care management are trained to help caregivers and people aged 60 years or older with behavioural and psychological symptoms of dementia.⁷ Psychological, physical, and educational interventions are combined to increase the effectiveness of these interventions in improving the quality of life of older people and their caregivers.

Some Hubs focus on integrating digital technology into scalable models of service delivery. In India, digital technology is used to train community health workers to deliver a brief evidence-based psychological treatment over six to eight sessions for depression in primary care settings in a relatively deprived state in the country.⁸ The low-cost training model replaces face-to-face training programmes requiring specialist trainers and costly logistical planning for delivery. In



John Naslund

Lancet Psychiatry 2020

Published Online

August 23, 2020

[https://doi.org/10.1016/](https://doi.org/10.1016/S2215-0366(20)30347-3)

[S2215-0366\(20\)30347-3](https://doi.org/10.1016/S2215-0366(20)30347-3)

Colombia, screening and treatment for depression and unhealthy alcohol use computerised validated clinical assessments, a digital clinical decision support tool to guide providers in treatment choices, and a digital therapeutic application to promote behaviour change and wellness among patients.⁹ In Mozambique, a brief electronic mental wellness tool helps community health workers, primary care providers, and mental health specialists to identify any mental disorder with accuracy. Additionally, the tool helps to provide standardised evidence-based interventions for severe and common mental disorders, alcohol and substance misuse, and suicide prevention through the use of digital therapeutic applications that are able to connect with electronic medical records.¹⁰

The Hubs rapidly responded to the COVID-19 crisis by adapting the above programmes to provide essential mental health and psychosocial support. Examples include developing modules to train teachers who are managing pandemic-related anxiety in school children in Pakistan, as well as taking measures to protect their own wellbeing. Technological innovations facilitating mental health delivery in Colombia have been mobilised and made freely available to assist with recovery efforts; furthermore, the Mozambican Ministry of Health has adopted and scaled up the technology as the new standard of care throughout the mental health system. In India, guidelines have been issued on the responsible reporting of suicides and other mental health concerns during COVID-19. In Malawi, a short telephone-based protocol is being used to identify suicide risk in community clinics, given that direct face-to-face contact is not possible. In India, Colombia, and Mozambique, digital tools are being deployed to manage stress, anxiety, and depression among frontline workers responding to patients with COVID-19.

Hubs present opportunities for mutual learning, in which frugal innovations forged out of necessity in LMICs can address racial and ethnic mental health disparities in high-income countries. For example, implementation strategies that have been developed and tested in Sierra Leone⁵ are being adapted for a US-based initiative targeting mental health disparities in migrant communities by advancing new, family-based preventive interventions. These interventions are being given by refugees, for refugees, to reduce the risks of mental health problems associated

with displacement, resettlement, family social and economic stress, experiences of racism, and poor access to care.

Collectively, the Hubs facilitate multidirectional learning, research capacity development, and, importantly, the design of novel implementation studies to advance scientific knowledge by examining the best strategies for scaling up mental health services in diverse settings, with all age groups, while addressing efficiency and cost-effectiveness. Much can be learnt from these successful collaborations to sustainably address immediate priorities and long-term strategies for mental health implementation research in LMICs, as well as in high-income countries, during and beyond COVID-19.

All authors are all supported by U19 'Scale Up Hubs' grants from the National Institute of Mental Health, USA. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

**Atif Rahman, John A Naslund, Theresa S Betancourt, Candace J Black, Anant Bhan, William Byansi, Hongtu Chen, Bradley N Gaynes, Carlos Gomez Restrepo, Lídia Gouveia, Syed Usman Hamdani, Lisa A Marsch, Inge Petersen, Ozge Sensoy Bahar, Laura Shields-Zeeman, Fred Ssewamala, Milton L Wainberg*
atif.rahman@liverpool.ac.uk

Institute of Population Health Sciences, University of Liverpool, Liverpool L69 3BX, UK (AR, SUH); Department of Global Health and Social Medicine, Harvard Medical School, Boston, MA, USA (JAN); Boston College School of Social Work, Boston, MA, USA (TSB, CJB); Sangath, Bhopal, India (AB); Brown School, Washington University in St Louis, St Louis, MO, USA (WB, OSB, FS); Department of Psychiatry, Brigham and Women's Hospital, Boston, MA, USA (HC); University of North Carolina School of Medicine, Chapel Hill, NC, USA (BNG); Pontificia Universidad Javeriana, Bogotá, Colombia (CGR); Department of Mental Health, Ministry of Health, Maputo, Mozambique (LG); Geisel School of Medicine, Dartmouth College, Hanover, NH, USA (LAM); Centre for Rural Health, College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa (IP); Netherlands Institute for Mental Health and Addiction (Trimbos Institute), Utrecht, Netherlands (LS-Z); and Department of Psychiatry, New York State, Columbia University, New York, NY, USA (MLW)

- 1 UN. Policy brief: COVID-19 and the need for action on mental health. Geneva: United Nations, 2020.
- 2 National Institute of Mental Health. Research partnerships for scaling up mental health interventions in low- and middle-income countries. 2020. <https://www.nimh.nih.gov/about/organization/cgmhr/scaleuphubs/index.shtml> (accessed July 9, 2020).
- 3 Hamdani SU, Muzaffar N, Huma Z, et al. 15-3 Using technology to advance school mental health: experience from the Eastern Mediterranean Region. *J Am Acad Child Adolesc Psychiatry* 2019; **58**: S22.
- 4 Ssewamala FM, Bahar OS, McKay MM, Hoagwood K, Huang KY, Pringle B. Strengthening mental health and research training in Sub-Saharan Africa (SMART Africa): Uganda study protocol. *Trials* 2018; **19**: 423.
- 5 Betancourt TS. Youth FORWARD: Scaling up an evidence-based mental health intervention in Sierra Leone. *Humanitarian Exchange* 2018; **72**: 30-32.
- 6 Pathare S, Shields-Zeeman L, Vijayakumar L, et al. Evaluation of the SPIRIT Integrated Suicide Prevention Programme: study protocol for a cluster-randomised controlled trial in rural Gujarat, India. *Trials* 2020; **21**: 1-3.

-
- 7 Chen H, Chuengsatiansup K, Levkoff S, et al. Using implementation science to advance management of dementia-related behavioral problems in a developing country. Alzheimer's Association International Conference 2018; Chicago, IL, USA; July 22–26, 2018; Chicago, 2018 (abstr P4-359).
 - 8 Muke SS, Shrivastava RD, Mitchell L, et al. Acceptability and feasibility of digital technology for training community health workers to deliver evidence-based psychosocial treatment for depression in rural India. *Asian Journal of Psychiatry* 2019; **45**: 99–106.
 - 9 Torrey WC, Cepeda M, Castro S, et al. Implementing technology-supported care for depression and alcohol use disorder in primary care in Colombia: preliminary findings. *Psychiatric Serv* 2020; **71**: 678–83.
 - 10 Wainberg ML, Lovero KL, Duarte CC, et al. Partnerships in research to implement and disseminate sustainable and scalable evidence based practices in Sub-Saharan Africa—Mozambique scale-up study protocol (PRIDE SSA-Mozambique). *Psychiatric Serv* (in press).